

## REFERENCES

- Willan R: Description and treatment of cutaneous diseases (order I). J Johnson London, 1798
- Plenck JJ: Doctrina de morbis cutaneis. Graeffer, Vienna, 1776
- Cazenave (PL)A, Schedel HE: Abrégé pratique des maladies de la peau Béchét Jeune, 3rd ed. Paris, 1838
- Bateman T: Delineations of cutaneous diseases: exhibiting the characteristic appearances of the principal genera and species comprised in the classification of the late Dr. Willan; and completing the series of engravings begun by that author. Longman, Hurst, Rees, Orme, and Brown, London, 1817
- Hebra F: Acute Exantheme und Hautkrankheiten. In: Virchow (ed.). Handbuch der speciellen Pathologie und Therapie, Vol. III. F Enke, Erlangen, 1860
- Lorry C-A: Tractatus de morbis cutaneis. Cavelier, Paris, 1777
- Booth CC: Robert Willan, MD FRS, FSA. Br J Dermatol 80:459-468, 1968
- Booth CC: Robert Willan and his kinsmen. Med History 25:181-196, 1981
- Booth CC: The Fothergillian medals of the Medical Society of London. J R Coll Phys 15:254-258, 1981
- Abraham JJ: The two Fothergills (The 170th Oration Medical Society of London). Trans Med Soc (Lond) LXIV:276-291, 1946
- The Fothergillian Gold Medalists: Transactions of the Medical Society of London, Vol. 88. Milner & Sons, London, 1972

## 8-Methoxypsoralen Acts on Lymphocyte Membranes in the Dark

To the Editor:

With great interest we have read the article by Dr. Malinin et al, "Ultrastructural modifications of the plasma membrane in HUT 102 lymphoblasts by long-wave ultraviolet light, psoralen, and PUVA" (J Invest Dermatol 95:97-103, 1990). The authors described therein membrane alterations after incubation of the cells with a suprapharmacologic concentration of the photosensitizer 8-methoxypsoralen (8-MOP) in the dark and suggest that cell membrane components are the primary target in the 8-MOP action mechanism.

The dark effect of 8-MOP on lymphocyte surface structures has been well documented by ourselves and others in several investigations using different methods with regard to their membrane-related evidence. Among other things, it has been shown that, following high dosages of 8-MOP, parts of the glycocalyx [1] and the cytoplasm [2] were released from leucocytes and lymphocytes, respectively, into the medium under cell culture conditions. However, morphologic changes could not be observed under the influence of 8-MOP in therapeutic concentrations [3]. On the other hand, without any UV-irradiation 8-MOP was able to increase the membrane-associated cAMP levels in mononuclear leucocytes [4], possibly by inhibition of phosphodiesterase [5] after only 1 min incubation time. Furthermore 8-MOP has shown to inhibit both the PHA and Con A-induced proliferation of normal human peripheral blood lymphocytes in a time- and dose-dependent manner [6,7]. This was explained by the decrease in IL-2 receptor expression on PHA-stimulated lymphocytes [6]. As we have shown, the PHA-triggered cytokine release (macrophage slowing factor [8]) and the HLA-DR expression [7] were depressed by 8-MOP, too. In addition, 8-MOP was able to temporarily reduce the binding of sheep erythrocytes to lymphocytes, which is known as membrane interaction [7]. Finally, 8-MOP reduced the number of Pan T cells evaluated by specific monoclonal antibodies [9].

Taken together, 8-MOP exerts some dark effects on lymphocyte surface membranes. These results may suggest some immunoregulatory effects of 8-MOP on (antigen activated) cells. Because of their only slight degree and transient nature [4,6-9], these 8-MOP dark effects are probably of low clinical relevance under therapeutic conditions.

Wolfgang Gast  
Uwe-Frithjof Hausteine  
Department of Dermatology  
Karl Marx University  
Leipzig, Germany

## REFERENCES

- Schmidt H, Weigel C: Zur Wirkung von Methoxalen auf menschliche Lymphozyten. Dermatol Monatsschr 174:20-27, 1988
- Roeder B, Krylenkov VA, Potapenko AY, Meffert H: Modifikation der Lymphozytenglykokalyx und der E-Rosetten-Bildung durch 8-MOP und UVA. Dermatol Monatsschr 169:157-165, 1983
- Schmoeckel C, Scherer R, Kern B, Braun-Falco O: The cytolytic effect of PUVA treatment on PHA-stimulated human peripheral lymphocytes. Acta Dermato Venerol 58:203-211, 1978
- Albrightson CR, Fertel RE, Brown BV, Stephens R: Psoralens increase the concentration of cyclic AMP in human cells in vitro. J Invest Dermatol 85:264-268, 1985
- Roemer R, Kittler L, Loeber G: Inhibitory effects of furocoumarins plus near UV light (365 nm) on cAMP phosphodiesterase activity. Stud biophys 94:33-36, 1983
- Cox GW, Orosz CG, Fertel RH: 8-methoxypsoralen inhibits lymphocyte proliferation in vitro in the absence of ultraviolet radiation. Int J Immunopharmacol 9:475-481, 1987
- Gast W, Rytter M, Walther T, Schaller J, Hofmann C, Hausteine UF: Dark effects of 8-methoxypsoralen on human lymphocytes. Stud biophys 124:173-178, 1988
- Rytter M, Gast W, Walther T, Herold W, Hausteine UF: 8-methoxypsoralen inhibits the release of macrophage slowing factor from lymphocytes. Stud biophys 133:109-112, 1989
- Schaller J, Gast W, Walther T, Hausteine UF, Hofmann C, Rytter M: Dark effects of 8-methoxypsoralen on lymphocyte surface markers. Photodermatology 4:310-311, 1987

## REPLY

In their letter concerning our article [1], Drs. Gast and Hausteine note, firstly, that "dark effect of 8-MOP on lymphocyte surface structures is well documented," and secondly conclude that "because of their only slight degree and transient nature these 8-MOP dark effects are probably of low clinical relevance."

We agree with the authors that dark 8-MOP reaction with lymphocytes and, with other cells, was indeed documented [2]; however, the availability of diverse phenomenologic data does not automatically imply understanding of underlying reaction mechanisms, and their biologic significance. Because photoreactions of 8-MOP, at least in part, are preceded and facilitated by dark reactions with specific targets, their ultimate significance cannot be extrapolated solely on the basis of their transience or detection by a single method.